

Yusif A. MAMMADOV, Tahir S. GADJIEV, Rafiq A. RASULOV

ON BEHAVIOR OF SOLUTIONS OF NONLINEAR PARABOLIC EQUATIONS

Abstract

The aim of this paper is to study the behavior of solutions of initial boundary problem for nonlinear parabolic equation with boundary regime. From physical point of view the rule of a boundary regime, the speed is more of movement in the arising temperature of a wave.

1. Introduction

In this paper the unbounded increasing solution of the nonlinear parabolic type equation for the finite times is considered. These type equations describe the processes of electrical and ionic heat conductivity in plasma, diffusion of neutrons and α -particles and etc. Investigation of unbounded solution of regime of peaking solutions occurs in the theory of nonlinear equations where one the essential ideas is the representation called eigen-function of nonlinear dissipative surroundings. It is well known that even a simple nonlinearity, subject to critical of exponent, the solution of nonlinear parabolic type equation for the finite time may increase unboundedly, i.e., there is a number $T > 0$ such that

$$\|u(x, t)\|_{L_\infty(R^n)} \rightarrow \infty, \quad t \rightarrow T < \infty.$$

In [1] the existence of unbounded solution for finite time with a simple nonlinearity has been proved. In [2] has been shown that any nonnegative solution subject to critical exponent is unbounded increasing for the finite time. Similar results were obtained in [3] an corresponding theorems are called Fujita-Hayakawa's theorems. More detailed reviews can be found in [4], [5] and [6].

2. Main results

Let us consider the equation

$$\frac{\partial u}{\partial t} = \sum_{i,j=1}^n \frac{d}{dx_j} \left(\left| \frac{\partial u}{\partial x_i} \right|^{p-2} \frac{\partial u}{\partial x_i} \right) - F(x, t, u) \quad (1)$$

In bounded domain $\Omega \subset R^n, n \geq 2$, with no smooth boundary, and mainly the boundary $\partial\Omega$ contains the conical points with span of the corner $\omega \in (0, \pi)$. Denote

$$\begin{aligned} \Pi_{a,b} &= \{(x, t) | x \in \Omega, a < t < b\}, \quad \Pi_a = \Pi_{a,\infty}, \\ \Gamma_{a,b} &= \{(x, t) | x \in \partial\Omega, a < t < b\}, \quad \Gamma_a = \Gamma_{a,\infty}. \end{aligned}$$

The functions $F(x, t, u), \frac{\partial F(x, t, u)}{\partial u}$ are continuous with respect to u uniformly in $\bar{\Pi}_0 \times \{u : |u| \leq M\}$ for any $M < \infty, F(x, t, 0) \equiv 0, \frac{\partial F}{\partial u}|_{u=0} \equiv 0$. Besides, the function F is measurable on whole arguments and doesn't decrease with respect to u . Let the Dirichlet boundary condition

$$u = f(x, t), \quad x \in \partial\Omega \quad (2)$$

the initial condition

$$u|_{t=0} = \varphi(x) \tag{3}$$

fulfilled on some domain $\Pi_{0,a}$, where $\varphi(x)$ is a smooth function. The condition on $f(x, t)$ is following

$$f'_t \in L_1((0, T_0), L_{\sigma+1}(\Omega)) \tag{4}$$

$$f \in L_1((0, T_0) \times \Omega) \cap ((0, T_0) \times L_{\sigma+1}(\Omega)) \quad T_0 < T.$$

Furthermore,

$$f(x, t) \rightarrow \infty, \quad t \rightarrow T. \tag{5}$$

Condition (5) is called boundary regime with sharpening at the blow-up time T . As a solution we understand the generalized solution from the Sobolev space $W_m^1(\Pi_0, a')$ for all $a < a'$, and the solution of problem (1)-(3) either exists in Π_0 , or

$$\lim_{t \rightarrow T-0} \max_{\Omega} |u(x, t)| = +\infty \tag{6}$$

at some $T = const.$

Assume that

$$\sup u \in \{x \in R^n, |x| < d\}, \quad 1 < d < R \tag{7}$$

$$\sup f \in [0, T] \times B_r, \quad r < R \tag{8}$$

$$f(0, x) - \varphi(x) \in W_{p+1}^1(\Omega, \partial\Omega). \tag{9}$$

Here $W_{p+1}^1(\Omega, \partial\Omega)$ is a closing in the norm $W_{p+1}^1(\Omega)$ and the functions from $C^\infty(\Omega)$ vanishing near $\partial\Omega$. Let's formulate the auxiliary results from [10], [11]. Let

$$L_p u = \operatorname{div} \left(|\nabla u|^{p-2} \nabla u \right), \quad p > 1$$

where p is a harmonic operator.

Lemma1. [10] *There exists the positive eigen-value of spectral problem for the operator L_p , to which corresponds the positive in Ω eigen-function.*

Lemma2. [11] *Let $u, v \in W_p^1(\Omega)$ and $u \leq v$ on $\partial\Omega$,*

$$\int_{\Omega} L_p(u) \eta_{x_i} dx \leq \int_{\Omega} L_p(v) \eta_{x_i} dx$$

for any $W_{p+1}^1(\Omega)$ with $\eta \geq 0$. Then $u \leq v$ on whole domain Ω .

We will call the generalized solution of problem (1)-(3) in $\Pi_{a,b}$ $u(x, t) \in W_p^1(\Pi_{a,b}), u(x, t) \in L_\infty(\Pi_{a,b})$ such that

$$- \int_{\Pi_{a,b}} \frac{\partial u}{\partial t} \psi dx dt + \sum_{i,j=1}^n \int_{\Pi_{a,b}} \left| \frac{\partial u}{\partial x_i} \right|^{p-2} \frac{\partial u}{\partial x_i} \frac{\partial \psi}{\partial x_j} dx dt = \int_{\Pi_{a,b}} F(x, t, u) \psi(x, t) dx dt. \tag{10}$$

Where $\psi(x, t)$ is an arbitrary function from $W_p^1(\Pi_{a,b}), \psi|_{\Gamma_{a,b}} = 0, 0 < a < b$ are any numbers. Let $u_0(x) > 0$ be an eigen-function of spectral problem for the operator

L_p corresponding to $\lambda = \lambda_1 > 0, \int_{\Omega} u_0(x) dx = 1, \int_{\Omega} u_0^q(x) dx = 1, q \geq 1$. Let's assume that the condition

$$(L_p u_0, u) \geq (L_p u, u_0)$$

Is fulfilled, i.e.,

$$\int_{\Omega} \left(\left| \frac{\partial u_0}{\partial x_i} \right|^{p-2} - \left| \frac{\partial u}{\partial x_i} \right|^{p-2} \right) \frac{\partial u}{\partial x_i} \frac{\partial u_0}{\partial x_i} dx \geq 0.$$

Let's introduce the function which will be characterized by blow-up boundary regime at $t \rightarrow T$.

$$G(t) = \lambda_1 \int_{\Omega} u_0(x) f(x, t) dx + \int_{\Omega} \left| f'_t \right|^{\sigma+1} dx + \int_{\Omega} f^{\sigma+1} dx,$$

for $t < T$.

Theorem 1. Let $f(x, t, u) \geq \alpha_0 |u|^{\sigma-1} u$ at $(x, t) \in \Pi_0, u \geq 0, \sigma > 1, \alpha_0 = \text{const} > 0$. There exists $k = \text{const}$ such that, if $u(x, 0) \geq 0, \int_{\Omega} u(x, 0) u_0(x) dx \geq k$, conditions (4), (5) and $G(t) \rightarrow \infty, t \rightarrow T$ are fulfilled, then $\lim_{t \rightarrow T-0} \max_{\Omega} u(x, t) = \infty, T = \text{const} > 0$.

Proof. Assume the vice versa, then $u(x, t)$ is a solution of equation (1) in Π_0 and condition (2) is fulfilled on Γ_0 . By virtue of Lemma 2, by lemma 1.1.2 $u_0(x) > 0$ in Π_0 . Substitute in (10) $\psi(x, t) = \varepsilon^{-1}(u_0(x) - f(x, t)), b = a + \varepsilon, a > 0, \varepsilon > 0$, where $u_0(x) > 0$ in Ω is eigen- function of spectral problem for the operator L_p , corresponding to eigen-value $\lambda_1 > 0$. Such eigen-value and eigen-function exist by virtue of Lemma 1. Then we

$$\begin{aligned} & -\frac{\varepsilon^{-1}}{2} \left[\int_{\Omega} u_0(x) u(x, a + \varepsilon) dx - \int_{\Omega} u_0(x) u(x, a) dx \right] + \\ & + \varepsilon^{-1} \int_{\Pi_{a, a+\varepsilon}} \left| \frac{\partial u}{\partial x_i} \right|^{p-2} \frac{\partial u}{\partial x_i} \frac{\partial \psi}{\partial x_j} dx dt - \varepsilon^{-1} \int_{\Pi_{a, a+\varepsilon}} F(x, t, u) \psi(x, t) dx dt = 0. \end{aligned} \quad (11)$$

Using condition (A), as ε approaches to zero in equality (11), we have

$$\begin{aligned} -g'(t) \leq & -\lambda_1 \int_{\Omega} u_0(x) u(x, t) dx - \int_{\Omega} u \frac{\partial f(x, t)}{\partial t} dx + \lambda_1 \int_{\Omega} u_0(x) f(x, t) dx - \\ & - \int_{\Omega} u^{\sigma} f(x, t) dx + \int_{\Omega} u_0 u^{\sigma} dx, \end{aligned} \quad (12)$$

where $g(t) = \int_{\Omega} u_0(x) u(x, t) dx$.

At first, let's estimate the second and the fourth integral on the right in (12). Apply the Cauchy inequality with $\varepsilon > 0$ to get

$$\int_{\Omega} u \frac{\partial f(x, t)}{\partial t} dx = \frac{\varepsilon_1}{2} \int_{\Omega} u^2 dx + C_1(\varepsilon_1) \int_{\Omega} f_t'^2 dx \leq$$

$$\leq \frac{\varepsilon_1}{2} \left(\int_{\Omega} u^{\sigma+1}(x, t) dx \right) + C_1(\varepsilon_1) \int_{\Omega} f_t'^{\sigma+1} dx + C_3 m \varepsilon \Omega,$$

$$\int_{\Omega} u^{\sigma} f(x, t) dx \leq \varepsilon_1 \int_{\Omega} u^{\sigma+1} dx + C_3(\varepsilon_1) \int_{\Omega} f^{\sigma+1}(x, t) dx.$$

Now, let's estimate $\int_{\Omega} u^{\sigma+1} dx$ and $\int_{\Omega} u_0 u^{\sigma} dx$. By using the Holder inequality for these integrals under the normed conditions $u_0(x)$ we have

$$\int_{\Omega} u^{\sigma+1} dx \geq g^{\sigma+1}(t), \quad \int_{\Omega} u_0 u^{\sigma} dx \geq g^{\sigma}(t).$$

Substituting these estimates in (12) to get

$$g'(t) \geq -\lambda_1 g(t) + g^{\sigma+1}(t) + \lambda_1 \int_{\Omega} u_0(x) f(x, t) dx +$$

$$+ \int_{\Omega} (f_t')^{\sigma+1} dx + \int_{\Omega} f^{\sigma+1}(x, t) dx. \quad (13)$$

Then from (13) we obtain

$$g'(t) \geq -\lambda_1 g(t) + C g^{\sigma+1}(t) + G(t). \quad (14)$$

Note that depending on a boundary regime there are various physical processes. At the power boundary regime with aggravation there is a heat localization. Such boundary regimes describe the solutions of equation stopped and thermal wave. Position of a point in front of a wave doesn't change in a current of all time of an aggravation $t \in (0, T)$ and thermal indications from the localization due to the fact that the temperature infinitely increase every where at $t \rightarrow T$. At this any boundary regime with an aggravation provides localization of thermal influence. For example, if $G(t) \leq (T-t)^{-1/\sigma}$, $\sigma > 1$, $0 < t < T$, there is a localization of heat, and at $G(t) \leq (T-t)^n$, where $n < -1/\sigma$, there is no localization. If $g(0) > C_2 = \left(\frac{\lambda_1}{C}\right)^{1/\sigma+1}$, then from (14), under the condition on $G(t)$, we'll obtain $\lim_{t \rightarrow T-0} g(t) = +\infty$. Hence we have $\lim_{t \rightarrow T-0} \max_{\Omega} u(x, t) = \infty$.

Thus, if $u(x, 0) \geq 0$ is not enough small, then the solution of equation (1) has been in Π_0 . Let's show that if $|u(x, 0)|$ is sufficiently small then solution of problem (1), (2), (3) exists on whole domain Π_0 . Here an important role is played by (9) and the condition which will coordinate the speeds of "attinity" of boundary and initial functions, i.e., for existence of solution at unboundedly tendency of boundary function to ? it is necessary sufficiently, small initial function.

Theorem 2. Let $|F(x, t, u)| \leq (C_1 + C_2 t^m) |u|^{\sigma}$, $\sigma > 1$. Relative to the boundary function, the conditions of Theorem 1 is fulfilled and there exists $\delta > 0$ such that if $|\varphi(x)| \leq \delta$, then solution of problem (1)-(3) exists in Π_0 , and $|u(x, t)| \leq C e^{-\alpha t}$, $\alpha = \text{const} > 0$.

Proof. Let $\bar{\Omega} \subset B_R$, where $B_R = \{x : |x| \leq R\}$ and $v > 0 \in B_R$ be an eigenfunction corresponding to the eigen-value λ_1 of boundary-value problem

$$L_m u + \lambda u = 0, x \in \Omega, u = 0, x \in \partial\Omega. \tag{15}$$

Consider the function $V(x, t) = \varepsilon e^{-\lambda_1 t/2} v(x)$. We have

$$V_t - L_m V - F(x, t, V) = \frac{1}{2} \varepsilon \lambda_1 e^{-\frac{\lambda_1 t}{2}} v(x) - (C_1 + C_2) \varepsilon^\sigma e^{-\frac{\lambda_1 t}{2}} v^\sigma, (x, t) \in \Pi_0$$

$$V > 0, (x, t) \in \Gamma_0.$$

If $\varepsilon > 0$ is sufficiently small, inequality (16) is understood in weak sense. From (16) and Lemma 2 follows that $|u| \leq V \leq C e^{-\lambda_1 t/2}$, if $|\varphi(x)| \leq \delta = \varepsilon \min_{\bar{\Omega}} v(x)$. No

let's determine the class of functions $\omega(x, t)$ consisting at the functions continuous in $\bar{\Pi}_{-\infty, +\infty}$, being equal to zero at $t \leq T$, $|\omega| \leq K e^{-ht}$. K and h such that $\bar{\Pi}_{-\infty, +\infty}$, $\|\omega\| = \sup_{\Pi_{-\infty, +\infty}} |\omega e^{ht}|$ is subset at Banach space continuous in $\theta(t) \subset C^\infty(R^1)$, $\theta(t) = 0$ functions with the norm $t \leq \tau$. Let $\theta(t) = 1$ and $t > T + 1$, and let's determine the operator H and K substituting $H\omega = \theta(t)z, \omega \in K$. Thus, we solve the linear problem

$$\sum_{i,j=1}^n \frac{d}{dx_j} \left(\left| \frac{\partial z(x, t)}{\partial x_i} \right|^{m-2} \frac{\partial z}{\partial x_i} \right) \theta(t) \cdot \omega_{x_i x_j} \frac{\partial z}{\partial t} + \theta(t) F(x, t, v) = 0.$$

Instead of the nonlinear problem. $z = 0$ at $(x, t) \in \Gamma_{-\infty, +\infty}$, consisting in the determination $\omega(x, t)$ in Ω by the known function $z(x, t)$. By means of semi-linear estimates above H terms K in K if T is sufficiently big. The operator H is quite continuous. This appears from the obtained estimation and theorem on Hölderiness of $\Pi_{-a, a}$, at solutions of parabolic equations for any a (see [1]). From Lere-Shouder theorem, the operator H has fixed point z . Theorem is proved.

References

[1]. Ladyzhenskaya O.A., Uraltseva N.N., Solonnikov V.A. *Linear and quasilinear equations of parabolic type*. M., Nauka, 1967 (Russian)

[2]. Chanillo S., Wheeden R. *Weighted Poincare and Sobolev inequalities and estimates for weighted Peano maximal functions* // Amer.J. Math., 1985, vol.707, No 5, pp.1191-1226.

[3]. Deng K., Levine H. *The role of critical exponents in blow-up theorems* // j.Math. Anal. Appl. 2000, vol.243, pp.85-126.

[4]. Ebihara Y., Kawashima Sh., Levine N. "On solution $u_{tt} - |x|^\alpha \Delta u = f(u)$ ", // Funccial Ekvac, 1995, 38, No 3, pp.539-544.

[5]. Fujita. *On the blocking up of solutions to the Cauchy problem for $U_t = \Delta U + U^{1+\alpha}$* , J. Fac.Sci. Univ.Tokyo, I. 1966, 13, pp.109-124.

[6]. Fujita. *On the blocking up of solutions to the Cauchy problem for $U_t = \Delta U + U^{1+\alpha}$* , J.Fac.Sci. Univ.Tokyo, I. 1966, 13, pp.109-124.

[Yu.A.Mammadov, T.S.Gadjiev, R.A.Rasulov]

[7]. Galaktionov V.A., Levine H. *The role of critical Fujita exponents and systems* // Nonlinear Anal., 1998, vol.34, pp.1005-1027.

[8]. Gilbarg D., Trudinger N. *Elliptic partial differential equations with derivations of the second order*. Moscow, Nauka, 1989.

[9]. Hayakawa K.O. *Nonexistence of global solutions of some semi linear parabolic differential equations*, Proc. Japan Acad. Ser. A. Math. Sci., 1973, 49, pp.503-505.

[10]. Kaplan S. *On the Growth of solutions of quasilinear parabolic equations*. II Comm. Pure Appl. Math., 1963, vol.16, pp. 305-330.

[11]. Levine H. *"Instability and nonexistence of global solutions to nonlinear wave equations of the form $Pu_{tt} = -Au + F(u)$ "*. //Trans. Amer. Math. Soc., 1974, 192, pp.1-21.

[12]. Samarskii A.A., Galaktionov V.A. Kurdyumov, Mikahailov. *Blow-up in quasi linear parabolic equations*. Moskow, Nauka, 1987.

[13]. Tolksdorf P. *On quasi linear boundary value problems in domains with corners*// Nonlinear Anal. 1981, vol. 5, No 7, pp.721-735.

Yusif A. Mammadov,

Azerbaijan State Pedagogical University
34, U. Hajibeyov str. Az 1000, Baku, Azernaijan

Tahir S. Gadjiev,

Institute of Mathematics and Mechanics of NAS of Azerbaijan
9, B.Vahabzade str., AZ1141, Baku, Azerbaijan
Tel.: (99412) 538 62 17 (off.).
tgadjiev@mail.az

Rafiq A. Rasulov

The Sheki Filial of the Institute of Teachers, Sheki,
Az5, Republic of Azerbaijan
rafig_rasulov@yahoo.com
Tel.: (99412) 539 47 20

Received: September 10, 2012; Revised: December 17, 2012.